

# 風治散

## THIS MEDICINE FOR GENERAL USE

Fuujisan Granules

(Category 2 OTC medicines)

## INDICATIONS :

Following symptoms in person with moderate or more physical strength; initial stage of common cold (without spontaneous sweating), coryza, rhinitis, headache, shoulder stiffness, muscle pain, and pain in hands and shoulder.

## CAUTION FOR USE

• Who should seek consultation

1. The following persons should consult a physician, pharmacist or registered distributor before taking this medicine.
  - (1) Persons undergoing medical treatment by a physician.
  - (2) Pregnant women or women who may be pregnant.
  - (3) Persons who is physically weak (people with reduced physical strength or a weak constitution).
  - (4) Persons with weak digestive systems.
  - (5) Persons with sweat excessively.
  - (6) Elderly persons.
  - (7) Persons who have experienced rash, redness, itching by medicines or other substances.
  - (8) Persons with the following symptoms: edema, difficulty in urination.
  - (9) Persons diagnosed as having the following: hypertension, heart disease, kidney disease, thyroid disease.
2. The following symptoms may be adverse reactions to this medication. If any of these symptoms occur, stop taking this medicine immediately and consult a physician, pharmacist or registered distributor. Take the instruction leaflet with you.

Affected Area	Symptoms
Skin	Rash, redness, itching
Gastrointestinal system	Nausea, loss of appetite, gastric distress

In very rare cases, the following serious symptoms may occur after taking medicine.  
In this case, consult a physician immediately.

Symptom name	Symptoms
Pseudoaldosteronism, myopathy	In addition to listlessness, numbness, a stretched feeling and stiffness in the hands and feet, feeling of weakness and muscular pain may appear and gradually worsen.
Hepatic function failure	Following symptoms appear: fever, itching, rash, jaundice (yellowing of skin and white of eyes), brown urine, whole-body tiredness, loss of appetite.

- Stop taking this medicine and consult a physician, pharmacist or registered distributor if your symptoms do not improve after taking this medicine for about 1 month (Take this medicine 5 to 6 times if you are in the early stages of a common cold, with symptoms such as a runny nose or headache). Take this leaflet with you.
- If you take this medicine regularly for a long period, consult a physician, pharmacist, or registered distributor.

## DOSAGE

Take the following doses with warm or cold water before or after meals.

Age	Dose	Daily Frequency
Adult (15 years or over)	1 Packet (2.5g)	3 times a day
7 years to under 15 years	2/3 Packet	
4 years to under 7 years	1/2 Packet	
2 years to under 4 years	1/3 Packet	

Less than 2 years	Do not use
-------------------	------------

#### CAUTION RELATED WITH DOSAGE AND USAGE

Children should take this medicine only under adult supervision.

#### INGREDIENTS

3 Packets (Adult daily dose)

Composition	Dosage	
Kakkon (Pueraria Root)	4.0g	2.8g of a water extract of the mixed crude drugs.
Maou (Ephedra Herb)	3.0g	
Shokyo (Ginger)	1.0g	
Taiso (Jujube)	3.0g	
Keihi (Cinnamon Bark)	2.0g	
Shakuyaku (Peony Root)	2.0g	
Kanzo (Glycyrrhiza)	2.0g	

Inactive ingredients: starch

#### CAUTION

Since the medicine contains crude drugs, Granule color may vary for between products, however this has no influence on the efficacy of this medicine.

#### PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Store in a cool, dry place free from direct sunlight.
- (2) Keep out of reach of children.
- (3) Do not transfer to another container to prevent misuse and quality deterioration.
- (4) When you take this medicine in divided doses, fold over the opened edge of the packet and store it. The remaining medicine should be taken within 2 days.

#### MANUFACTURED AND DISTRIBUTED BY:

WAKANYAKU MEDICAL INSTITUTE LTD.

1193 Akagisan, Fujimi-machi, Maebashi-shi, Gunma. Japan

Prepared: July 2025 (Third edition)