

ササイサン

THIS MEDICINE FOR GENERAL USE

Sasaisan

(Category 2 OTC medicines)

INDICATIONS :

Loss of appetite, stomach and abdominal bloating, dyspepsia, gastrointestinal weakness, overeating, overdrinking, heartburn, heavy stomach, the pressure on the chest, nausea (including retching, upset stomach, sick feeling by hangover and/or sick from drinking), vomiting

CAUTION FOR USE

• Who should seek consultation

1. The following persons should consult a physician, pharmacist or registered distributor before taking this medicine.
 - (1) Persons undergoing medical treatment by a physician.
 - (2) Elderly persons.
 - (3) Persons with the following symptom: edema.
 - (4) Persons diagnosed as having the following disease: hypertension, kidney disease, heart disease.
2. In very rare cases, the following serious symptoms may occur after taking medicine.
In this case, consult a physician immediately.

Symptom name	Symptoms
Pseudoaldosteronism, myopathy	In addition to listlessness, numbness, a stretched feeling and stiffness in the hands and feet, feeling of weakness and muscular pain may appear and gradually worsen.

3. Stop taking this medicine and consult a physician, pharmacist or registered distributor if your symptoms do not improve after taking this medicine for about 1 month of use.

Take this leaflet with you.

4. If you take this medicine regularly for a long period, consult a physician, pharmacist, or registered distributor.

DOSAGE

Take the following doses with warm or cold water, between meals.

(“Between meals” means a couple of hours after meals.)

Age	Dose	Daily Frequency
Adult (15 years or over)	1 Packet (1.5g)	3 times a day
11 years to under 15 years	2/3 Packet	
8 years to under 11 years	1/2 Packet	
5 years to under 8 years	1/3 Packet	
3 years to under 5 years	1/4 Packet	
Less than 3 years	Do not use	

CAUTION RELATED WITH DOSAGE AND USAGE

Children should take this medicine only under adult supervision.

INGREDIENTS

3 Packets (Adult daily dose)

Composition	Dosage
Kanzo (Glycyrrhiza) extract (Equivalent to 3.6g of raw Glycyrrhiza)	0.9g
Oubaku (Phellodendron Bark) extract (Equivalent to 2.5g of raw Phellodendron Bark)	0.4g
Hakka (Mentha Herb)	0.3g

Inactive ingredients: lactose hydrate

CAUTION

Since the medicine contains crude drugs, Granule color may vary for between products, however this has no influence on the efficacy of the medicine.

PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Store in a cool, dry place free from direct sunlight.
- (2) Keep out of reach of children.
- (3) Do not transfer to another container to prevent misuse and quality deterioration.
- (4) When you take this medicine in divided doses, fold over the opened edge of the packet and store it. The remaining medicine should be taken within 2 days.

MANUFACTURED AND DISTRIBUTED BY:

WAKANYAKU MEDICAL INSTITUTE LTD.

1193 Akagisan, Fujimi-machi, Maebashi-shi, Gunma. Japan

Prepared: July 2025 (Third edition)