

# 新ササカル (顆粒)

## THIS MEDICINE FOR GENERAL USE

Shin Sasa-cal Granules

(Category 3 OTC medicines)

## INDICATIONS :

a) Growth promotion of bones and/or teeth in the following cases; weak constitution, scrofulosis\*

\*Scrofulosis refers to a weakened constitution with a tendency to develop anemia.

b) Prevention of weakness of bones and/or teeth during pregnancy or lactation

## CAUTION FOR USE

• Who should seek consultation

1. The following persons should consult a physician, pharmacist or registered distributor before taking this medicine.

Persons undergoing medical treatment by a physician.

2. The following symptoms may be adverse reactions to this medication. If any of these symptoms occur, stop taking this medicine immediately and consult a physician, pharmacist or registered distributor. Take the instruction leaflet with you.

Affected Area	Symptoms
Skin	Rash, redness, itching

3. The following symptom may appear after taking this medicine. If the symptom continues or worsen, stop taking this medicine and consult a physician, pharmacist or registered distributor. Take this leaflet with you.: constipation

4. If you take this medicine regularly for a long period, consult a physician, pharmacist, or registered distributor.

## DOSAGE

Take the following doses with warm or cold water between meals.

(“Between meals” means a couple of hours after meals.)

Age	Dose	Daily Frequency
15 years or over	1 Packet (1.5g)	3 times a day
7 years to under 15 years	2/3 Packet	
5 years to under 7 years	1/2 Packet	

## CAUTION RELATED WITH DOSAGE AND USAGE

- (1) Follow the dosage and directions.
- (2) Children should take this medicine only under adult supervision.

## INGREDIENTS

3 Packets (Adult daily dose)

Composition	Dosage	Function
Borei powder (Powdered Oyster Shell)	1.5g (Containing 470mg of Calcium)	Calcium supplements
Calcium Gluconate Hydrate	0.8g (Containing 70mg of Calcium)	Calcium supplements
Calcium Lactate Hydrate	0.5g (Containing 60mg of Calcium)	Calcium supplements
Dried Yeast	0.3g	Supplements of B vitamins, amino acids, etc., and increasing calcium absorption

Inactive ingredients: starch, lactose hydrate, sodium copper chlorophyllin, flavor

## CAUTION

Since the medicine contains crude drugs, Granule color may vary for between products, however this has no influence on the efficacy of the medicine.

#### PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Store hermetically in a cool dry place free from direct sunlight.
- (2) Keep out of reach of children.
- (3) Do not transfer to another container to prevent misuse and quality deterioration.
- (4) When you take this medicine in divided doses, fold over the opened edge of the packet and store it. The remaining medicine should be taken within 2 days.

#### MANUFACTURED AND DISTRIBUTED BY:

WAKANYAKU MEDICAL INSTITUTE LTD.

1193 Akagisan, Fujimi-machi, Maebashi-shi, Gunma. Japan

Prepared: February 2026 (First edition)