

ササクール A

THIS MEDICINE FOR GENERAL USE

Sasa-cool A

(Category 2 OTC medicines)

INDICATIONS :

Hoarseness, Throat discomfort

CAUTION FOR USE

• Who should seek consultation

1. The following persons should consult a physician, pharmacist or registered distributor, before taking this medicine.
 - (1) Persons undergoing medical treatment by a physician.
 - (2) Pregnant women or women who may be pregnant.
 - (3) Persons with a weak stomach and intestines who tend to have diarrhea.
 - (4) Persons who have experienced rash, redness or itching, or similar reactions by medicines or other substances.
2. The following symptoms may be adverse reactions to this medication. If any of these symptoms occur, stop taking this medicine immediately and consult a physician, pharmacist or registered distributor. Take the instruction leaflet with you.

Affected Area	Symptoms
Skin	Rash, redness, itching
Gastrointestinal system	Loss of appetite, gastric distress

3. Stop taking this medicine and consult a physician, pharmacist or registered distributor if your symptoms do not improve after taking this medicine for 5 to 6 days. Take this leaflet with you.

CAUTION ASSOCIATED INDICATION

The medicine can be used regardless of physical strength.

DOSAGE

Place each dose in the mouth, allow it to dissolve slowly, and then swallow. Take between meals.

(“Between meals” means a couple of hours after meals.)

Age	Dose	Daily Frequency
Adult (15 years or over)	50 Pellets	3 times a day
7 years to under 15 years	30 Pellets	
4 years to under 7 years	25 Pellets	
2 years to under 4 years	12 Pellets	
Less than 2 years	Do not use	

CAUTION RELATED WITH DOSAGE AND USAGE

- (1) Follow the dosage and directions.
- (2) Children should take this medicine only under adult supervision.
- (3) When young children take this medication, be careful it does not get stuck in the throat.

INGREDIENTS

Composition	150 Pellets (Daily dosage)	
Rengyo (Forsythia Fruit)	0.75g	0.75g of a water extract of the mixed crude drugs
Kikyo (Platycodon Root)	0.75g	
Kanzo (Glycyrrhiza)	0.75g	
Shukusya (Amomum Seed)	0.30g	
Senkyu (Cnidium Rhizome)	0.30g	
Kashi (Myrobalan)	0.30g	

Asenyaku (Gambir)	0.60g	
Hakka (Mentha Herb)	1.20g	
Rengyo powder (Powdered Forsythia Fruit)	188mg	
Kikyo powder (Powdered Platycodon Root)	188mg	
Kanzo powder (Powdered Glycyrrhiza)	188mg	
Shukusya powder (Powdered Amomum Seed)	75mg	
Senkyu powder (Powdered Cnidium Rhizome)	75mg	
Kashi powder (Powdered Myrobalan)	75mg	
Asenyaku powder (Powdered Gambir)	150mg	
Hakka powder (Powdered Mentha Herb)	300mg	

Inactive ingredients: starch

CAUTION

Since the medicine contains crude drugs, Pellet color may vary for between products, however this has no influence on the efficacy of the medicine.

PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Store in a cool, dry place free from direct sunlight.
- (2) Keep out of reach of children.
- (3) Do not transfer to another container to prevent misuse and quality deterioration.
- (4) When you take this medicine in divided doses, fold over the opened edge of the packet and store it. The remaining medicine should be taken within 2 days.

MANUFACTURED AND DISTRIBUTED BY:

WAKANYAKU MEDICAL INSTITUTE- LTD.

1193 Akagisan, Fujimi-machi, Maebashi-shi, Gunma. Japan

Prepared: July 2025 (Third edition)