

和漢薬晴快散

THIS MEDICINE FOR GENERAL USE

Wakanyaku Seikaisan Granules

(Category 3 OTC medicines)

INDICATIONS :

For forgetfulness occurring in middle age and beyond

CAUTION FOR USE

• To be avoided;

Failure to observe the following may result in symptom worsening or adverse reactions.

The following persons do not use.

- (1) Persons diagnosed as having the following disease: dementia.
- (2) Persons suspected with dementia.

• Who should seek consultation

1. The following persons should consult a physician, pharmacist or registered distributor before taking this medicine.
 - (1) Persons undergoing medical treatment by a physician.
 - (2) Pregnant women or women who may be pregnant.
 - (3) Persons who have experienced allergic symptom by medicines or other substances.
2. In case of the following, stop taking this medicine immediately and consult a physician, pharmacist or registered distributor. Take the instruction leaflet with you.
 - (1) the following symptoms occur, after taking this medicine.

Affected Area	Symptoms
Skin	Rash, redness, itching
Gastrointestinal system	Nausea, vomiting, loss of appetite, gastric
Neuropsychiatric system	Dizziness

3. Stop taking this medicine and consult a physician, pharmacist or registered distributor if your symptoms do not improve after taking this medicine for about 1 month. Take this leaflet with you.
4. The following symptom may occur after taking medicine. If the continuance or enhancement of such a symptom is observed, stop taking this medicine and consult a physician, pharmacist or registered distributor. Take this leaflet with you.
: Diarrhea
5. If you take this medicine regularly for a long period, consult a physician, pharmacist, or registered distributor.

OTHER CAUTIONS

Taking this medicine may influence the laboratory value (1,5-AG in the blood) regarding diabetes.

DOSAGE

Take the following doses with warm or cold water before or between meals.

(“Between meals” means a couple of hours after meals.)

Age	Dose	Daily Frequency
Adult (15 years or over)	1 Packet (1.2g)	3 times a day
Less than 15 years	Do not use	

INGREDIENTS

3 Packets (Adult daily dose)

Composition	Dosage
Onji (Polygara Root) extract (Equivalent to 3g of raw Polygara Root)	1.3g

Inactive ingredients: dextrin, magnesium stearate, light anhydrous silicic acid, crystalline cellulose, lactose hydrate

CAUTION

Since the medicine contains crude drugs, Granule color may vary for between products, however this has no influence on the efficacy of this medicine.

PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Store in a cool, dry place free from direct sunlight.
- (2) Keep out of reach of children.
- (3) Do not transfer to another container to prevent misuse and quality deterioration.
- (4) When you take this medicine in divided doses, fold over the opened edge of the packet and store it. The remaining medicine should be taken within 2 days.

MANUFACTURED AND DISTRIBUTED BY:

WAKANYAKU MEDICAL INSTITUTE LTD.

1193 Akagisan, Fujimi-machi, Maebashi-shi, Gunma. Japan

Prepared: July 2025 (Third edition)