

紫華榮

THIS MEDICINE FOR GENERAL USE

Shikaron

(Category 3 OTC medicines)

INDICATIONS

For revitalization in individuals with the following conditions: Weak constitution, physical fatigue, weakness during and after illness, gastrointestinal weakness, loss of appetite, pale complexion, over-sensitive to cold, and revitalization in growth season.

CAUTION FOR USE

• Who should seek consultation

1. The following symptoms may be adverse reactions to this medication. If any of these symptoms occur, stop taking this medicine immediately and consult a physician, pharmacist or registered distributor. Take the instruction leaflet with you.

Affected Area	Symptoms
Skin	Rash, redness, itching

2. If you take this medicine regularly for a long period, consult a physician, pharmacist, or registered distributor.

DOSAGE

Take the following doses with warm or cold water between meals.

(“Between meals” means a couple of hours after meals.)

Age	Dose	Daily Frequency
Adult (15 years or over)	1 Packet (1.5g)	3 times a day
11 years to under 15 years	1/2 Packet	
8 years to under 11 years	1/3 Packet	

CAUTION RELATED WITH DOSAGE AND USAGE

- (1) Observe the dosage and usage.
- (2) Children should take this medicine only under adult supervision.

INGREDIENT

3 Packets (Adult daily dose)

Composition	Raw material	Dosage
Shikon (Lithospermum Root) extract	7,500mg	170mg
Touki (Japanese Angelica Root) extract	342mg	120mg
Jyuyaku (Houttuynia Herb) extract	1,500mg	85mg
Ninjin (Ginseng) extract	3,000mg	600mg
Ougi (Astragalus Root) extract	250mg	50mg
Senkyu (Cnidium Rhizome) extract	225mg	50mg
Yokuinin powder (Powdered Coix Seed)		1,800mg
Kanzo powder (Powdered Glycyrrhiza)		45mg

Inactive ingredients: lactose hydrate

CAUTION

Since the medicine contains crude drugs, Granule color may vary for between products, however this has no influence on the efficacy of the medicine.

PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Store in a cool dry place free from direct sunlight.
- (2) Keep out of reach of children.
- (3) Do not transfer to another container to prevent misuse and quality deterioration.
- (4) When you take this medicine in divided doses, fold over the opened edge of the packet and store it. The remaining medicine should be taken within 2 days.

MANUFACTURED AND DISTRIBUTED BY:

WAKANYAKU MEDICAL INSTITUTE LTD.

1193 Akagisan, Fujimi-machi, Maebashi-shi, Gunma. Japan

Prepared: July 2025 (Third edition)