

松寿仙

THIS MEDICINE FOR GENERAL USE

Shojusen

(Category 3 OTC medicines)

INDICATIONS :

For revitalization in individuals with the following conditions; weak constitution, physical fatigue, during and after illness, gastrointestinal weakness, loss of appetite.

CAUTION FOR USE

• Who should seek consultation

1. The following persons should consult a physician, pharmacist or registered distributor before taking this medicine.
 - (1) Persons undergoing medical treatment by a physician.
 - (2) Persons who are allergic in person or family.
 - (3) Persons who have experienced allergic symptoms by medicines or something.
2. The following symptoms may be adverse reactions to this medication. If any of these symptoms occur, stop taking this medicine immediately and consult a physician, pharmacist or registered distributor. Take the instruction leaflet with you.

Affected Area	Symptoms
Skin	Rash, redness, itching

3. If you take this medicine regularly for a long period, consult a physician, pharmacist, or registered distributor.

DOSAGE

Take the following doses measured with the attached measuring cup, after dilution with warm or cold water, before or between meals.

(“Between meals” means a couple of hours after meals.)

Age	Dose	Daily Frequency
Adult (15 years or over)	2ml	3 times a day
8 years to under 15 years	1ml	

CAUTION RELATED WITH DOSAGE AND USAGE

- (1) Follow the dosage and directions.
- (2) Children should take this medicine only under adult supervision.

INGREDIENT

Composition (in 6ml)	Dosage (in 6ml)
Chlorophyllous liquid of Sasa (Equivalent to 19g of raw Sasa)	4.8ml
Akamatsu-ba extract (Japanese red pine needles extract) (Equivalent to 232mg of raw Japanese red pine needles)	18.0mg
Ninjin extract (Ginseng extract) (Equivalent to 58mg of raw Ginseng)	10.8mg

Inactive ingredients: paraben

CAUTION

Since the medicine contains crude drugs, Liquid color may vary for between products, however this has no influence on the efficacy of the medicine.

PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Keep out of reach of children.
- (2) Store in a cool place free from direct sunlight.
- (3) Not to touch the spout directly when opening and taking to prevent quality deterioration by a contamination from hands.
- (4) Be careful not to cut fingers when opening and separating off a plastic loop.
- (5) Store in refrigerator with putting cap back on the bottle after opening, and use up as early as possible.

- (6) After opening, content fluid may spill in case of carrying the bottle with the part of cap and/or keeping the bottle sideways under a condition of failing to close the cap tightly.
- (7) Do not transfer to another container to prevent misuse and quality deterioration.
- (8) Since the medicine is herbal extract formulation, ingredient such as amino acid may attach to the side or the bottom of the bottle, however this has no influence on the efficacy of the medicine by taking after shaking the bottle.

MANUFACTURED AND DISTRIBUTED BY:

WAKANYAKU MEDICAL INSTITUTE LTD.

1193 Akagisan, Fujimi-machi, Maebashi-shi, Gunma. Japan

Prepared: July 2025 (Third edition)